Draft version February 12, 2010
Preprint typeset using LaTeX style emulateapj v. 08/22/09

arXiv:1002.2534v1 [astro-ph.CO] 12 Feb 2010

INTERMEDIATE AGE STARS AS ORIGIN OF THE LOW VELOCITY DISPERSION NUCLEAR RING IN MRK 1066

Rogemar A. Riffel[1,2], Thaisa Storchi-Bergmann[2], Rogério Riffel[2] and Miriani G. Pastoriza[2]
[1]Universidade Federal de Santa Maria, Departamento de Física, Centro de Ciências Naturais e Exatas, 97105-900, Santa Maria, RS, Brazil
[2]Universidade Federal do Rio Grande do Sul, IF, CP 15051, 91501-970, Porto Alegre, RS, Brazil

Draft version February 12, 2010

ABSTRACT

We report the first two-dimensional stellar population synthesis in the near-infrared of the nuclear region of an active galaxy, namely Mrk 1066. We have used integral field spectroscopy with adaptative optics at the Gemini North Telescope to map the to map the age distribution of the stellar population in the inner 300 pc at a spatial resolution of 35 pc. An old stellar population component (age $\gtrsim$5 Gyr) is dominant within the inner $\approx$160 pc, which we attribute to the galaxy bulge. Beyond this region, up to the borders of the observation field ($\sim$300 pc), intermediate age components (0.3–0.7 Gyr) dominate. We find a spatial correlation between this intermediate age component and a partial ring of low stellar velocity dispersions (σ_*). Low-σ_* nuclear rings have been observed in other active galaxies and our result for Mrk 1066 suggests that they are formed by intermediate age stars. This age is consistent with an origin for the low-σ_* rings in a past event which triggered an inflow of gas and formed stars which still keep the colder kinematics (as compared to that of the bulge) of the gas from which they have formed. At the nucleus proper we detect, in addition, two unresolved components: a compact infrared source, consistent with an origin in hot dust with mass $\approx 1.9 \times 10^{-2}$ M$_\odot$, and a blue featureless power-law continuum, which contributes with only $\approx$15% of the flux at 2.12 μm.

Subject headings: galaxies: individual (Mrk 1066) — galaxies: nuclei — galaxies: Seyfert — galaxies: stellar content — infrared: galaxies

1. INTRODUCTION

Optical spectroscopy on scales of hundreds of parsecs around the nucleus of Seyfert galaxies have shown that in $\approx$40 % of them the active galactic nucleus (AGN) and young stars co-exist (e.g. Storchi-Bergmann et al. 2000, 2001; González Delgado, Heckman & Leitherer 2001; Cid Fernandes et al. 2004; Asari et al. 2007; Dors et al. 2008), providing support to the so-called AGN-Starburst connection (e.g. Norman & Scoville 1988; Terlevich, Diaz & Terlevich 1990; Heckman et al. 1997; Heckman 2004; Riffel et al. 2009a). In particular, these authors have pointed out that the main difference between the stellar population (hereafter SP) of active and non-active galaxies is an excess of mainly intermediate age stars in the former.

A similar result has been found in recent SP studies in the near-infrared (hereafter near-IR), using the technique of spectral synthesis (Riffel et al. 2009b, 2007). Using integrated spectra of the central few hundreds of parsecs in a sample of 24 Seyfert galaxies, these authors have shown that the continuum is dominated by the contribution of intermediate-age stellar population components (SPCs). In addition, they found that the near-IR nuclear spectra of about 50 % of the Seyfert 1 and ∼20 % of the Seyfert 2 galaxies show emission from hot dust (Riffel et al. 2009b; Riffel, Storchi-Bergmann & McGregor 2009; Riffel et al. 2009a; Rodríguez-Ardila & Mazzalay 2006; Rodríguez-Ardila, Contini & Viegas 2005a).

Using a somewhat distinct technique, Davies et al. (2007) obtained near-IR integral field spectroscopy to investigate the circumnuclear star formation in 9 nearby Seyfert galaxies at spatial resolutions of tens of parsecs. They have modelled the Brγ equivalent width, supernova rate and mass-to-light ratio to quantify the star formation history in the center of these galaxies using their code STARS. They found that the ages of the stars which contribute most to the near-IR continuum lie in the range 10–30 Myr, but point out that these ages should be considered only as “characteristic”, as they have not performed a proper spectral synthesis, arguing that there may be simultaneously two or more SPs that are not coeval (Davies et al. 2007, 2006).

In this paper we present, for the first time, two-dimensional (hereafter 2D) SP synthesis in the near-IR for the inner hundreds of pc of an active galaxy – namely Mrk 1066 – using integral field spectroscopy with adaptive optics, which allowed us to derive the contribution of distinct SPCs to the near-IR spectra and map their spatial distributions. This paper is organized as follows. In Sec. 2 we describe the observations, data reduction procedures and the spectral synthesis method; in Sec. 3 we present our results, which are discussed in Sec. 4. The conclusions are presented in Sec. 5.

2. THE DATA AND SYNTHESIS CODE

2.1. *Mrk 1066*

Mrk 1066 is an SB0 galaxy harboring a Seyfert 2 nucleus and located at a distance of 48.6 Mpc, for which 1″corresponds to 235 pc at the galaxy. Previous studies of the central region of Mrk 1066 have shown that line emission from high-excitation gas is dominated by gas in a bi-conical outflow oriented along position angle PA=135/315°, which seems to be associated to the radio jet, while the low-excitation gas is more restricted to the plane of the galaxy (Bower et al. 1995; Nagar et al. 1999; Knop et al. 2001; Riffel, Storchi-Bergmann & Nagar 2010; Riffel & Storchi-Bergmann 2010). We selected this galaxy for the present study because it shows a signature of the presence of young/intermediate age stellar population in its stellar kinematics in the inner few hundred parsecs as illustrated in the stellar velocity dispersion (σ_*) map shown in the top-left panel of Fig. 2. This map was obtained from the fit of the

Electronic address: rogemar@smail.ufsm.br

CO band heads around 2.3 μm by a combination of stellar spectral templates from Winge, Riffel & Storchi-Bergmann (2009) using the penalized Pixel-Fitting (pPXF) method of Cappellari & Emsellem (2004). It shows a partial ring of low σ_*, a signature of the presence of stars which still have the kinematics of the cold gas from which they have formed[1]. Our first goal with the spectral synthesis is to derive the age of this population. Additionally, high spatial resolution spectroscopy has shown that the near-IR nuclear continuum is consistent with a large contribution from hot dust emission (Riffel, Storchi-Bergmann & Nagar 2010), not detected in lower resolution observations (Riffel et al. 2009b). Our second goal is to quantify this contribution with spectral synthesis.

2.2. *Observations and data reduction*

Mrk 1066 was observed with the Gemini Near-Infrared Integral-Field Spectrograph (NIFS; McGregor et al. 2003) operating with Gemini North Adaptive Optics system ALTAIR in September 2008 under the programme GN-2008B-Q-30. Two set of observations were obtained, one at the J band, covering a spectral region from 1.15 μm to 1.36 μm with a two-pixel spectral resolving power of 6040; and another at the K_l band at resolving power of 5290 and covering the spectral range 2.1–2.5 μm.

The data reduction, described in Riffel, Storchi-Bergmann & Nagar (2010), was performed using the GEMINI IRAF package and followed standard procedures. The resulting calibrated data cube contains 784 spectra covering a angular region of $2''\!.8\times2''\!.8$ at a sampling of $0''\!.1\times0''\!.1$, which corresponds to $650\times650\,\mathrm{pc}^2$ and $23.5\times23.5\,\mathrm{pc}^2$, respectively. The angular resolution is $\approx0''\!.15$, corresponding to 35 pc at the galaxy.

2.3. *Analysis*

In order to model the continuum and stellar absorption features we used the STARLIGHT spectral synthesis code (Cid Fernandes et al. 2004, 2005, 2009; Mateus et al. 2006; Asari et al. 2007). According to the authors STARLIGHT mixes computational techniques originally developed for semi empirical population synthesis with ingredients from evolutionary synthesis models.

In summary, the code fits an observed spectum O_λ with a combination, in different proportions, of N_* single stellar populations (SSPs). Following Riffel et al. (2009b) we use the Maraston (2005) Evolutionary Population Synthesis (EPS) models as base set (described below). This choice is due to the fact that Maraston's models include the effects of stars in the TP-AGB phase, wich is crucial to model NIR SP (see Riffel, Rodríguez-Ardila & Pastoriza 2006; Riffel et al. 2007, 2008b, 2009b). Extinction is modeled by STARLIGHT as due to foreground dust, and parameterized by the V-band extinction A_V (Cid Fernandes et al. 2004, 2005). Note that all the components are reddened by the same ammount. We use the CCM (Cardelli, Clayton & Mathis 1989) extinction law.

Basically, the code solves the following equation for a model spectrum M_λ (Cid Fernandes et al. 2005):

$$M_\lambda = M_{\lambda 0}\left[\sum_{j=1}^{N_*} x_j\, b_{j,\lambda}\, r_\lambda\right] \otimes G(v_*, \sigma_*) \qquad (1)$$

were $b_{j,\lambda}\, r_\lambda$ is the reddened spectrum of the jth SSP normalized at λ_0; $r_\lambda = 10^{-0.4(A_\lambda - A_{\lambda 0})}$ is the reddening term; $M_{\lambda 0}$ is the synthetic flux at the normalisation wavelength; $\vec{x}$ is the population vector; $\otimes$ denotes the convolution operator and $G(v_*, \sigma_*)$ is the gaussian distribution used to model the line-of-sight stellar motions, it is centred at velocity v_* with dispersion σ_*.

The final fit is carried out with a simulated annealing plus Metropolis scheme, which searches for the minimum of the equation (Cid Fernandes et al. 2005):

$$\chi^2 = \sum_\lambda [(O_\lambda - M_\lambda) w_\lambda]^2 \qquad (2)$$

where emission lines and spurious features are masked out by fixing w_λ=0.

The spectral base was constructed with EPS models of Maraston (2005) as in Riffel et al. (2009b), and comprise single stellar populations (SSPs) synthetic spectra covering 12 ages (t =0.01,0.03, 0.05, 0.1, 0.3, 0.5, 0.7, 1, 2, 5, 9 and 13 Gyr) and four metallicities (Z =0.02, 0.5, 1, 2 $Z_\odot$). We also included blackbody functions for temperatures in the range 700-1400 K in steps of 100 K (Riffel et al. 2009b) and a power-law ($F_\nu \propto \nu^{-1.5}$) in order to account for possible contributions from dust emission and from a featureless continuum (FC), respectively, in regions close to the nucleus (e.g. Cid Fernandes et al. 2004).

3. RESULTS

In Fig. 1 we present the results of the synthesis for two spectra extracted within an aperture of $0''\!.1\times0''\!.1$, one at the nucleus (top panel) and the other at 1$''$ south-east of it (bottom panel). The observed (black) and synthetic (red) spectra were normalized to unit at 2.12 μm. This wavelength was chosen because the K-band spectra present a higher signal-to-noise ratio than those in the J-band and the spectral region near 2.12 μm is free of emission/absorption lines (Riffel et al. 2008b). In the inner $0''\!.2$ (45 pc) radius the contribution from a FC and from dust emission are important, as illustrated in the top panel of Fig. 1, where the combined contribution of all SPCs is shown in blue, the FC as a dotted line and the composite blackbody function (sum over all temperatures) as a dashed line. We point out that the syntetic spectra (sum of all components) was reddened by the reddening amount detected in the observed spectra.

Following Cid Fernandes et al. (2004), we have binned the contribution of the SPCs (x_j) to the flux at 2.2μm into four age (t) ranges: young (x_y: $t \leq 100$ Myr); young-intermediate (x_{yi}: $0.3 \leq t \leq 0.7$ Gyr), intermediate-old (x_{io}: $1 \leq t \leq 2$ Gyr) and old (x_o: $5 \leq t \leq 13$ Gyr). The percent flux contribution at 2.12 μm from stars of each age range are shown in Fig. 2: on average, the young-intermediate SPCs dominate the continuum emission from the central region of Mrk 1066, but there is significant variation over the NIFS field. While in regions farther than $r\ \sim0''\!.7$ – 160 pc from the nucleus – the contribution of the young-intermediate age stars reaches values of up to 100 %, closer to the nucleus its contribution is negligible. Within this region the stellar population is dominated by the old component, whose contribution reaches $\approx$ 50 %, followed by the young component, whose contribution reaches $\approx$30 % of the flux at 2.12 μm. Also within this region, but unresolved by our observations (thus within $0''\!.15$ from the nucleus) we find a contribution of up to 50% from the combined blackbody components. Finally, we have found a small contribution from the AGN FC component, which reaches at most

---

[1] A detailed discussion of the stellar kinematics is presented in Riffel & Storchi-Bergmann (2010).



Fig. 1.— Sample fits of the SPCs to the nuclear spectrum (top) and to an extra-nuclear one at 1.″0 south-east of the nucleus (bottom). The observed spectra are shown in black and the fits in red. In the top panel the dashed line shows the contribution of hot dust, the dotted line that of the featureless continuum and the blue continuous line shows the contribution of the combined SPCs.

15 % at the nucleus, also unresolved by our observations. The intermediate-old component contributes with less than 20 % at most locations, with larger values more concentrated to the west-northwest of the nucleus.

Besides the SPC distributions, the synthesis outputs the average reddening of the stellar populations, shown in the central panel of Fig. 3. It reaches the highest values, of up to $E(B-V) = 1.4$, along the position angle 128°, which is the orientation of the line of nodes of the galaxy (Riffel & Storchi-Bergmann 2010).

The robustness of the SP fit can be measured by the percent mean deviation (*adev*), $|O_\lambda - M_\lambda|/O_\lambda$, where O_λ is the observed spectrum and M_λ is the fitted model (Cid Fernandes et al. 2004, 2005). The resulting *adev* map is shown in the right panel of Fig. 3 and presents values $adev \lesssim 2.5\,\%$ at most locations, indicating that the model reproduces very well the observed spectra. Nevertheless, at regions close to the border of the NIFS field the *adev* reaches values of up to 5%. At these locations we found an increase in the contribution of the young component x_y, which has thus larger uncertainties than in the rest of the NIFS field and should be further investigated with better data.

4. DISCUSSION

In general, our results are in good agreement with those of previous near-IR studies for single aperture nuclear spectra (Riffel et al. 2009b; Ramos Almeida, Pérez García & Acosta-Pulido 2009). The main novelty of our work is the 2D mapping of the stellar population using near-IR spectra, which shows spatial variations in the contribution of the SPCs in the inner few hundred parsecs of Mrk 1066.

The significance of the above variations is further enhanced by the comparison with the σ_* map presented in the top-left panel of Fig. 2. This map shows a partial ring of low σ_* values ($\approx 50\,\mathrm{km\,s^{-1}}$) surrounding the nuclear region (which has $\sigma_* \approx 100\,\mathrm{km\,s^{-1}}$) at $\approx 1''$ from it. Such rings are commonly observed in the central region of active galaxies (Barbosa et al. 2006; Deo, Crenshaw & Kraemer 2006; Simões Lopes et al. 2007; Riffel et al. 2008a, 2009a) and interpreted as being due to colder regions with more recent star formation than the underlying bulge. The comparison of the results from the SP synthesis of Mrk 1066 with the σ_* map shows that the low σ_*-values ring is associated with the young-intermediate age SPC, while the highest σ_* values are associated with the old SPC, supporting the use of low stellar velocity dispersion as a tracer of younger stars in the bulge and confirming the interpretation of the above studies.

The flux-weighted SPC contributions depend on the normalization point and thus the comparison with results from other spectral regions should be done with caution. A physical parameter which does not depend on the normalization point and spectral range used in the synthesis is the mass of the stars. Thus, we constructed maps for the mass-weighted contribution of each SPC. The contributions for the young population (m_y), young-intermediate population (m_{yi}), intermediate-old population (m_{io}) and old population (m_o), are shown in the rightmost panel of second row, bottom-left, bottom-central and bottom-right panels of Fig. 2, respectively. The mass-weighted contribution of the young population is very small over the whole field, while m_o dominates within $\approx$160 pc from the nucleus and m_{yi} dominates in the circumnuclear ring.

The only previous 2D stellar population studies of active galaxies in the near-IR published to date are those from the group of R. I. Davies. In particular, Davies et al. (2007) investigated the circumnuclear star formation in nine Seyfert galaxies using near-IR IFU observations and also found circumnuclear disks of typical diameters of tens of pc with a "characteristic" age in the range 10–300 Myr based on measurementes of the Brγ emission-line equivalent width, supernova rate and mass-to-light ratio. Thus, the results we have found for Mrk 1066 are in reasonable agreement with those found by Davies group. Nevertheless, the methodology adopted in the



FIG. 2.— Stellar velocity dispersion map (top-left panel) and spatial distributions of the percent contribution of each SPC to the flux at λ = 2.12μm (x_j) and to the mass (m_j), where j represents the age of the SPC: young (y: ≤ 100 Myr), young-intermediate (yi: 0.3–0.7 Gyr), intermediate-old (io: 1–2 Gyr) and old (o: 5–13 Gyr).



FIG. 3.— From left to right: Spatial distribution of the dust emission summed over all temperatures, reddening map and *adev* map (percent mean deviation from the spectral fit).

present work allowed us not only to obtain a "characteristic" age, but to also map, for the first time, the spatial distribution of stars of different ages in the central region of a Seyfert galaxy, on the basis of near-IR integral-field spectroscopy.

As presented in Sec. 2.3 our spectral base include four distinc metalicities: Z =0.02, 0.5, 1, 2 $Z_\odot$. Following Cid Fernandes et al. (2004), we can estimate the mean flux and mass-weighted metalicity in the fit by

$$< Z_* >_L = \sum_{j=1}^{N_*} x_j Z_j \tag{3}$$

and

$$< Z_* >_M = \sum_{j=1}^{N_*} m_j Z_j , \tag{4}$$

where $< Z_* >_L$ is the flux-weighted mean metalicity, $< Z_* >_M$ is the mass-weighted mean metalicity and N_* is the number of SSP in the spectral base. In the case of Mrk 1066 we obtain a nearly solar metalicity – $< Z_* >_L = 0.021$ and $< Z_* >_M = 0.017$ – in good agreement with those found by Riffel et al. (2009b) from near-IR spectral synthesis using a single aperture nuclear spectrum.

In further support of the results of the synthesis, we found that the average reddening map derived for the stellar population (central panel of Fig. 3) is in close agreement to the one derived for the narrow-line region using emission-line ratios (Riffel, Storchi-Bergmann & Nagar 2010), presenting a similar S-shaped structure.

The synthesis also confirmed the presence of an unresolved blackbody (BB) component at the nucleus, which we had found in Riffel, Storchi-Bergmann & Nagar (2010) from the fit of the nuclear spectrum by a BB function plus a power-law. On the basis of its luminosity, we have followed the calculations as in Riffel et al. (2009b) to estimate a total dust mass of $M_{\rm HD} \approx 1.9 \times 10^{-2}$ M$_\odot$, which is in good agreement with the value obtained in Riffel, Storchi-Bergmann & Nagar (2010) and with those observed for other Seyfert galaxies (e.g. Riffel et al. 2009a; Riffel, Storchi-Bergmann & McGregor 2009; Riffel et al. 2009b; Rodríguez-Ardila & Mazzalay 2006; Rodríguez-Ardila, Contini & Viegas 2005a).

As discussed in Sec. 3, we found that the FC component contributes with $\sim$ 15 % of the 2.12 μm nuclear continuum, which is in good agreement with the result found by Riffel et al. (2009b). This contribution is nevertheless uncertain, as it is hard to distinguish a reddened young starburst ($t \lesssim 5$ Myr) from an AGN FC component (e.g. Storchi-Bergmann et al. 2000; Cid Fernandes et al. 2004; Riffel et al. 2009b). On the other hand, the contribution of the young population x_y is dominated by stars with ages of 50–100 Myr, thus there does not seem to be a significant contribution from very young stellar populations, suggesting that the FC component originates from the emission of the AGN. This interpretation is supported by the detection of weak broad components for the Paβ and Brγ emission lines for the nuclear spectra of Mrk 1066 (Riffel, Storchi-Bergmann & Nagar 2010; Veilleux,Goodrich & Hill 1997). However, spectropolarimetric studies, show no evidence for the hidden Seyfert 1 nucleus in Mrk 1066 (e.g. Bian & Gu 2007). Thus, the origin of the FC in Mrk 1066 must be further investigated.

We finally point out one caveat of the stellar population synthesis in the near-IR. In Riffel, Storchi-Bergmann & Nagar (2010), on the basis of emission-line ratios, we have identified a large star-forming region at $0\farcs5$ south-east of the nucleus. In order to be active, this region should have an age $\lesssim$10 Myr (Riffel et al. 2009a; Dors et al. 2008; Díaz et al. 2007; Kennicutt et al. 1989). Nevertheless, our synthesis did not find a young stellar component at this location, indicating that spectral synthesis in the near-IR (at least in the J and K bands) is not a good tracer of very recent star formation (age $\leq$ 10 Myr). This result is expected since the young (blue) population has its peak emission in ultraviolet/optical wavelengths, while in the near-IR its contribution to the flux is very small, being hard to detect this component in this wavelength region.

In order to have a complete census of the stellar population components it is thus essential to study the stellar content of the central region of active galaxies using distinct methods and spectral regions.

5. CONCLUSIONS

The present work reports for the first time spectral synthesis in the near-IR with 2D coverage for the nuclear region of a Seyfert galaxy (Mrk 1066) within the inner $\approx$ 300 pc at a spatial resolution of $\approx$ 35 pc. We have mapped the distribution of stellar population components of different ages and of their average reddening. The main conclusions of this work are:

- The age of the dominant stellar population presents spatial variations: the flux and mass contributions within the inner $\approx$ 160 pc are dominated by old stars ($t \geq 5$ Gyr), while intermediate age stars ($0.3 \leq t \leq 0.7$ Gyr) dominate in the circumnuclear region;

- There is a spatial correlation between the distribution of the intermediate age component and low stellar velocity dispersion values which delineate a partial ring around the nucleus. Similar structures have been found around other active nuclei, and our result for Mrk 1066 suggests that these nuclear rings (and in some cases disks) are formed by intermediate age stars.

- There is an unresolved dusty structure at the nucleus with mass $M_{\rm HD} \approx 1.9 \times 10^{-2}$ M$_\odot$, which may be the hottest part of the dusty torus postulated by the unified model of AGN and a small contribution from a power-law continuum ($\approx$15 % of the flux at 2.12 μm);

- The near-IR synthesis seems not to be sensitive to very recent star formation (with $t \lesssim 5$ Myr), reinforcing the importance of multi-wavelength stellar population studies of the central region of active galaxies.

Based on observations obtained at the Gemini Observatory, which is operated by the Association of Universities for Research in Astronomy, Inc., under a cooperative agreement with the NSF on behalf of the Gemini partnership: the National Science Foundation (United States), the Science and Technology Facilities Council (United Kingdom), the National Research Council (Canada), CONICYT (Chile), the Australian Research Council (Australia), Ministério da Ciência e Tecnologia (Brazil) and south-eastCYT (Argentina). This work has been partially supported by the Brazilian institutions CNPq and CAPES.

REFERENCES

Asari, N. V., Cid Fernandes, R., Stasińska, G., Torres-Papaqui, J. P., Mateus, A., Sodré, L., Schoenell, W. & Gomes, J. M., 2007, MNRAS, 381, 263.
Barbosa, F. K. B., Storchi-Bergmann, T., Cid Fernandes, R., Winge, C., & Schmitt, H. 2006, MNRAS, 371, 170.
Bian, W. & Gu, C.. 2007, ApJ, 657, 159.
Bower, G., Wilson, A., Morse, J. A., Gelderman, R., Whitle, M., & Mulchaey, J., 1995, ApJ, 454, 106.
Cappellari, M., Emsellem, E. 2004, PASP, 116, 138.
Cardelli,J. A., Clayton, G. C. & Mathis, J. S., 1989, ApJ, 345,245.
Cid Fernandes, R., Gu, Q. Melnick, J., Terlevich, E., Terlevich, R., Kunth, D., Rodrigues Lacerda, R., Joguet, B., 2004, MNRAS, 355, 273.
Cid Fernandes, R., Mateus, A., Sodré, L., Stasińska, G., Gomes, J. M., 2005, MNRAS, 358, 363.
Cid Fernandes, R. et al. 2009, RMXAC, 35, 127. 2005, MNRAS, 358, 363.
Davies, R. I. et al. ApJ, 646, 754
Davies, R. I., Sánchez, F. M., Genzel, R., Tacconi, L. J., Hicks, E. K. S., Friedrich, S., & Sternberg, A. ApJ, 671, 1388
Deo, R. P., Crenshaw, D. M. & Kraemer, S. B., 2006, AJ, 132, 321.
Díaz, A. I., Terlevich, E., Castellanos, M., Hägele, G. F., 2007, MNRAS, 382, 251.
Dors, O. L., Storchi-Bergmann, T., Riffel, Rogemar A., & Schmidt, A., 2008, MNRAS
Ganda, K. et al., 2007, MNRAS, 380, 506.
González Delgado, R., Heckman, T., Leitherer, C., 2001, ApJ, 546, 845.
Heckman, T. M., González Delgado, R. M., Leitherer, C., Meurer, G. R., Krolik, J., Wilson, A. S., Koratkar, A., Kinney, A. 1997, ApJ, 482, 114.
Heckman, T. M., in Ho L. ed, Co-evolution of Black Holes and Galaxies, from the Carnegie Observatories Centennial Symposia. Carnegie Observatories Astrophysics Series. Cambridge University Press, Cambridge, pg. 358.
Kennicutt, R. C., Jr., Keel, W. C., Blaha, C. A., 1989, AJ, 97, 1022.
Knop, R. A., Armus, L., Matthews, K., Murphy, T. W., & Soifer, B. T., 2001, ApJ, 122, 764.
Maraston, C., 2005, MNRAS, 362, 799.
Mateus, A., Sodré, L., Cid Fernandes, R., Stasińska, G., Schoenell, W., Gomes, J. M., 2006, MNRAS, 370, 721
McGregor, P. J. et al., 2003, Proceedings of the SPIE, 4841, 1581.
Nagar, N. M., Wilson, A. S., Mulchaey, J. S. & Gallimore, J. F., 1999, ApJS, 120, 209.
Norman, C., & Scoville, N., 1988, ApJ, 332, 124.
Peletier, R. F. et al., 2007, MNRAS, 379, 445.
Raimann, D., Storchi-Bergmann, T., González Delgado, R. M., Cid Fernandes, R., Heckman, T., Leitherer, C., & Schmitt, H. 2003, MNRAS, 339, 792.
Ramos Almeida, C., Pérez García, A. M., & Acosta-Pulido, J. A., 2009, ApJ, 694, 1379.
Rodríguez-Ardila, A., Contini, M., & Viegas, S. M., 2005, MNRAS, 357, 220.
Rodríguez-Ardila, A., & Mazzalay, X., 2006, MNRAS, 367, 57.
Riffel, Rogemar A., Storchi-Bergmann, T., Winge, C., McGregor, P. J., Beck, T., Schmitt, H. 2008a, MNRAS, 385, 1129.
Riffel, Rogemar A., Storchi-Bergmann, T., Dors, O. L., & Winge, C., 2009a, MNRAS, 393, 783.
Riffel, Rogemar A., Storchi-Bergmann, T., McGregor, P. J. 2009, ApJ, 698, 1767.
Riffel, Rogemar A., Storchi-Bergmann, T., & Nagar, N., 2010, accepted by MNRAS, arxiv.org/abs/1001.0130.
Riffel, Rogemar A., & Storchi-Bergmann, T., 2010, submitted to MNRAS
Riffel, R., Rodríguez-Ardila, A. & Pastoriza, M. G., 2006, A&A, 457, 61.
Riffel, R., Pastoriza, M. G., Rodríguez-Ardila, A., & Maraston, C., 2007, ApJ, 659, 103.
Riffel, R., Pastoriza, M. G., Rodriguez-Ardila, A., Maraston, C., 2008b, MNRAS, 388, 803.
Riffel, R., Pastoriza, M. G., Rodriguez-Ardila, A., Bonatto, C. J., 2009, MNRAS, 400, 273.
Simões Lopes, R. D., Storchi-Bergmann, T., de Fátima Saraiva, M., & Martini, P., 2007, ApJ, 655, 718.
Storchi-Bergmann, T., Raimann, D., Bica, E. L. D., Fraquelli, H., A. 2000, ApJ, 544, 747.
Storchi-Bergmann, T., González Delgado, R., Schmitt, H., Cid Fernandes, R., Heckman, T., 2001, ApJ, 559, 147.
Terlevich, E., Diaz, A. I., & Terlevich, R. 1990, MNRAS, 242, 271.
Veilleux, S., Goodrich, R. W. & Hill, G. J., 1997, ApJ, 477, 631.
Winge, C., Storchi-Bergmann, T., Riffel, Rogemar A., 2009, ApJS, 185, 186.